

# UC San Diego

## UC San Diego Previously Published Works

### Title

Outcomes of Randomized Clinical Trials of Interventions to Enhance Social, Emotional, and Spiritual Components of Wisdom: A Systematic Review and Meta-analysis.

### Permalink

<https://escholarship.org/uc/item/5rt6n0gp>

### Journal

JAMA psychiatry, 77(9)

### ISSN

2168-622X

### Authors

Lee, Ellen E  
Bangen, Katherine J  
Avanzino, Julie A  
et al.

### Publication Date

2020-09-01

### DOI

10.1001/jamapsychiatry.2020.0821

Peer reviewed

# Outcomes of Randomized Clinical Trials of Interventions to Enhance Social, Emotional, and Spiritual Components of Wisdom

## A Systematic Review and Meta-analysis

Ellen E. Lee, MD; Katherine J. Bangen, PhD; Julie A. Avanzino, BA; BaiChun Hou, BS; Marina Ramsey; Graham Eglit, PhD; Jinyuan Liu, M.S.; Xin M. Tu, PhD; Martin Paulus, MD; Dilip V. Jeste, MD

**IMPORTANCE** Wisdom is a neurobiological personality trait made up of specific components, including prosocial behaviors, emotional regulation, and spirituality. It is associated with greater well-being and happiness.

**OBJECTIVE** To evaluate the effectiveness of interventions to enhance individual components of wisdom.

**DATA SOURCES** MEDLINE and PsycINFO databases were searched for articles published through December 31, 2018.

**STUDY ELIGIBILITY CRITERIA** Randomized clinical trials that sought to enhance a component of wisdom, used published measures to assess that component, were published in English, had a minimum sample size of 40 participants, and presented data that enabled computation of effect sizes were included in this meta-analysis.

**DATA EXTRACTION AND SYNTHESIS** Random-effect models were used to calculate pooled standardized mean differences (SMDs) for each wisdom component and random-effects meta-regression to assess heterogeneity of studies.

**MAIN OUTCOMES AND MEASURES** Improvement in wisdom component using published measures.

**RESULTS** Fifty-seven studies (N = 7096 participants) met review criteria: 29 for prosocial behaviors, 13 for emotional regulation, and 15 for spirituality. Study samples included people with psychiatric or physical illnesses and from the community. Of the studies, 27 (47%) reported significant improvement with medium to large effect sizes. Meta-analysis revealed significant pooled SMDs for prosocial behaviors (23 studies; pooled SMD, 0.43 [95% CI, 0.22-0.63];  $P = .02$ ), emotional regulation (12 studies; pooled SMD, 0.67 [95% CI, 0.21-1.12];  $P = .004$ ), and spirituality (12 studies; pooled SMD, 1.00 [95% CI, 0.41-1.60];  $P = .001$ ). Heterogeneity of studies was considerable for all wisdom components. Publication bias was present for prosocial behavior and emotional regulation studies; after adjusting for it, the pooled SMD for prosocial behavior remained significant (SMD, 0.4 [95% CI, 0.16-0.78];  $P = .003$ ). Meta-regression analysis found that effect sizes did not vary by wisdom component, although for trials on prosocial behaviors, large effect sizes were associated with older mean participant age ( $\beta$ , 0.08 [SE, 0.04]), and the reverse was true for spirituality trials ( $\beta$ , -0.13 [SE, 0.04]). For spirituality interventions, higher-quality trials had larger effect sizes ( $\beta$ , 4.17 [SE, 1.07]), although the reverse was true for prosocial behavior trials ( $\beta$ , -0.91 [SE 0.44]).

**CONCLUSIONS AND RELEVANCE** Interventions to enhance spirituality, emotional regulation, and prosocial behaviors are effective in a proportion of people with mental or physical illnesses and from the community. The modern behavioral epidemics of loneliness, suicide, and opioid abuse point to a growing need for wisdom-enhancing interventions to promote individual and societal well-being.

JAMA Psychiatry. doi:10.1001/jamapsychiatry.2020.0821  
Published online May 13, 2020.

[+ Editorial](#)

[+ Supplemental content](#)

**Author Affiliations:** Author affiliations are listed at the end of this article.

**Corresponding Author:** Dilip V. Jeste, MD, Sam and Rose Stein Institute for Research on Aging, Department of Psychiatry, University of California, San Diego, 9500 Gilman Dr, #0664, La Jolla, CA 92023 (djeste@ucsd.edu).

Wisdom has been discussed in religious and philosophical texts for centuries. The empirical study of wisdom began only 4 decades ago, but the number of research articles on wisdom has been growing. Vaillant's Harvard Study of Adult Development was the first to examine psychosocial and lifestyle factors associated with wisdom and well-being.<sup>1</sup> Baltes and Staudinger<sup>2</sup> defined wisdom as extensive pragmatic knowledge, focusing on the cognitive or intellectual aspect. Clayton and Birren<sup>3</sup> added reflective and affective components to the definition of wisdom. Sternberg<sup>4</sup> posited that wisdom resulted from an application of knowledge mediated by a balance of personal and societal interests. Ardel<sup>5</sup> conceptualized wisdom as an integration of cognitive, reflective, and affective (or compassionate) personality qualities. Cloninger's research<sup>6</sup> highlighted the relevance of certain personality traits to well-being.

We previously reviewed the empirical literature on wisdom<sup>7,8</sup> and conducted a Delphi method study of consensus among international experts in wisdom,<sup>7</sup> as well as a mixed-methods qualitative-quantitative study of wisdom in an ancient religious document. These studies suggested that wisdom is a complex human trait with several specific components (ie, prosocial behaviors such as empathy and compassion, emotional regulation, spirituality, self-reflection, social decision-making or social advising, acceptance of uncertainty, and decisiveness). Wisdom is thought to be adaptive rather than fixed and can increase with age and personal experience. Blazer,<sup>8</sup> Williams et al,<sup>9</sup> Grossman,<sup>10</sup> and others have emphasized practical wisdom, considering the context that influences wise decision-making. Several investigations have reported that wisdom is associated with positive outcomes, including better overall physical and mental health,<sup>11</sup> well-being,<sup>12</sup> happiness,<sup>13</sup> life satisfaction,<sup>14</sup> and resilience,<sup>15</sup> as well as lower levels of loneliness.<sup>15</sup>

We also reviewed the literature on neurobiological basis of wisdom components and found that they seemed to share similar brain regions, specifically the prefrontal cortex (dorsolateral, ventromedial, and anterior cingulate) and limbic striatum.<sup>16</sup> Furthermore, we found a number of published case reports of damage to these areas that produced a loss of these components (eg, increased emotional lability, impulsivity, indecisiveness, lack of compassion) without affecting other cognitive abilities, such as in the case of Phineas Gage<sup>17</sup> and patients with frontotemporal dementia.<sup>18</sup> While there have been no prospective longitudinal studies of changes in wisdom with aging using standardized rating scales, older adults have been reported in multiple investigations to have greater emotional regulation,<sup>19</sup> positivity,<sup>20</sup> prosocial behaviors,<sup>21</sup> and self-reflection<sup>19</sup> compared with younger adults. Considerable empirical evidence indicating neuroplasticity of aging, especially in adults who are physically, cognitively, and socially active, supports the potential to modify wisdom-type traits in later life. Aging-associated brain adaptations, including reduced lateralization of functioning,<sup>22</sup> posterior-to-anterior shift in brain activity,<sup>23</sup> and diminished amygdala response to negative or stressful stimuli,<sup>24</sup> may facilitate wisdom-relevant brain function in later life. Distinct patterns of brain activation on functional magnetic resonance imaging (fMRI) in response to

## Key Points

**Question** How effective are interventions to enhance individual components of wisdom?

**Findings** Despite heterogeneity of studies and publication bias, this meta-analysis and meta-regression found that interventions to enhance prosocial behaviors, emotional regulation, and spirituality were generally effective, especially among older participants.

**Meaning** Per this analysis, interventions to enhance prosocial behaviors, emotional regulation, and spirituality are effective in a proportion of individuals with mental or physical illnesses and people from the community.

moral dilemmas have been observed in individuals with higher vs lower scores on a validated scale for assessing overall wisdom.<sup>25</sup>

Personality traits, such as resilience and optimism, have been shown to be moderately heritable (with estimates of 33%<sup>26</sup> to 52%<sup>27</sup>), suggesting they are also influenced by environmental factors. A recent investigation<sup>28</sup> reported 50% to 58% heritability of self-directedness, cooperativeness, and self-transcendence in a large, population-based study that was replicated in multiple samples. These components overlap with those in our definition of wisdom. Several recent studies have reported increases in resilience and optimism with behavioral interventions.<sup>29,30</sup> While acknowledging their limitations, these findings support the possibility that positive personality traits can be enhanced through psychosocial interventions and thereby potentially lead to improvement of health and well-being. Yet we found no published reviews of interventions for increasing wisdom or its components; thus, it was unclear whether such interventions were effective. Therefore, we conducted a systematic review with meta-analysis and meta-regression analysis of randomized clinical trials (RCTs) intended to enhance one of the specific components of wisdom.

## Methods

### Procedures

We conducted a literature search for interventions targeting wisdom components, as outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) diagram (Figure 1). To identify articles for review, we surveyed the MEDLINE and PsycINFO online databases through December 31, 2018, with the following criteria: any study that (1) included one of the wisdom components listed above, (2) was published in English, (3) was an RCT, (4) had a minimum sample size of 40 participants, (5) included a published assessment tool to measure that component before and after the intervention, and (6) presented data that enabled computation of the intervention's effect size for enhancing the wisdom component.

The following MEDLINE search terms were used: “((Intervention [Title/Abstract] OR Interventions [Title/Abstract])

AND (wisdom[Title] OR pro-social[Title] OR empathy[Title] OR compassion[Title] OR emotion regulation[Title] OR emotional regulation[Title] OR spirituality[Title] OR self-reflection[Title] OR decisiveness[Title] OR social advising[Title] OR altruism[Title]) AND ("0001/01/01"[PDat]: "2018/12/31"[PDat]) AND English[lang]." The following search terms were used in the PsycINFO database: "(ti(interventions OR intervention) OR ab(interventions OR intervention)) AND ti(wisdom OR pro-social OR empathy OR compassion OR self-reflection OR emotion regulation OR emotional regulation OR spirituality OR decisiveness OR social advising OR altruism))," with the additional limits of English language and publication prior to December 31, 2018.

This search yielded 513 articles of potential interest after the removal of duplicates, of which 153 were deemed relevant based on a review of their abstracts. At least 2 authors (of a group of 4: E.E.L., J.A.A., B.H., and G.E.) independently examined each full-text journal article for defined eligibility criteria. The  $\kappa$  statistic for study selection was 0.97. Disagreements on inclusion criteria were settled by a third author from the same group of 4. The final search resulted in 57 studies reported in 54 articles.

### Statistical Analysis


For the meta-analysis, outcome data (assessment of wisdom component) were extracted from each study, including means, SDs, and sample sizes for the intervention and control groups. Data extraction was conducted by several authors in duplicate (E.E.L., M.R., and B.H.). The standardized mean differences (SMDs) for each RCT and the pooled SMD for each wisdom component were calculated using a random-effects model, given the heterogeneity of the interventions and outcome measures. Pooled SMDs were interpreted as small ( $0.2 \leq \text{SMD} < 0.5$ ), medium ( $0.5 \leq \text{SMD} < 0.8$ ), and large ( $0.8 \leq \text{SMD}$ ).<sup>31</sup>

The quality of these studies was evaluated using a modified Scale for Assessing Scientific Quality of Investigations (SASQI).<sup>32,33</sup> Thirteen original items were retained in the modified version; the 3 excluded items referred to educational interventions that used technology.

The  $I^2$  statistic was used to assess statistical heterogeneity of the study outcomes, with the following interpretation about its importance: low (0%-40%), moderate (30%-60%), substantial (50%-90%), and high (75%-100%).<sup>34</sup> Publication bias was assessed using the Egger test (a measure of the asymmetry of the funnel plot),<sup>35</sup> as well as the Duval and Tweedie trim-and-fill procedure<sup>36</sup> to adjust the funnel plot and pooled SMD for missing studies.

To perform meta-regression analyses, we first converted different types of effect sizes into  $\eta^2$  and logit-transformed  $\eta^2$  to remove range restriction. We then performed a meta-regression model using the generalized estimating equation, weighted by sample size of each study.<sup>37</sup> This led to models with the following factors included: type of patient (community based vs physically ill vs psychiatrically ill), mean age of participants, percentage of female participants, intervention format (individual vs group), number of sessions, length of individual sessions (minutes), and SASQI score (with

Figure 1. PRISMA Flow Diagram for Literature Review



the median split into 2 levels). The variance inflation factor was calculated to detect potential multicollinearity.<sup>38</sup> Additional factors included interaction terms: mean age by wisdom component, SASQI median split by wisdom component, and length of sessions by wisdom component based on variance inflation factor analyses. Significance was defined as an  $\alpha$  less than .05 (2-tailed) for all analyses. All analyses were conducted using RevMan version 5.3 (Cochrane Reviews) and R version 3.6.1 (R Foundation for Statistical Computing).

## Results

### Study Characteristics

Of the 57 intervention studies that met our review criteria (Table 1; eTable 1 in the Supplement), 29 focused on prosocial behaviors<sup>39-65</sup> (eg, empathy, compassion, altruism), 13 on emotional regulation,<sup>66-77</sup> and 15 on spirituality.<sup>78-92</sup> We did not find any published RCTs for self-reflection, social decision-making or social advising, acceptance of uncertainty, and decisiveness.

Forty-five studies<sup>43-52,55-60,62-65,67,70,71,73,74,76-92</sup> included only adult participants. Twenty-nine reports<sup>39,41-46,51,53,54,56-67,70,72,79,89,92</sup> had community-based participants, 19 reports<sup>40,47-49,52,55,68,69,71,73-78,81,90,91</sup> included persons with psychiatric illnesses or behavioral problems, and 9 had participants with physical illnesses.<sup>50,80,82-88</sup> Forty interventions<sup>39,42-44,46,49,50,53,54,56-62,64-68,71-76,80,81,83,84,86,87,89-92</sup> took place in a group setting. Across the wisdom components, the interventions varied in length, duration, and outcome measures used (eTable 1 in the Supplement). The mean (SD) intervention duration was 748 (753) minutes. Study quality was generally high because of the criteria for selecting reports for this review, with mean (SD) modified SASQI scores of 10.44 (1.7; possible range, 0-13; observed range, 5-13; eTable 2 in the Supplement). Compared with control

Table 1. Summary of Intervention Characteristics<sup>a</sup>

Characteristic	Wisdom component, No.		
	Prosocial behaviors	Emotional regulation	Spirituality
Randomized clinical trial			
With inert control group	17	8	8
With active control group	9	5	7
Age groups			
Child/adolescent	6	5	0
Adult	20	8	15 <sup>a</sup>
Sample characteristics			
Community based	18	5	3
Physically ill	1	0	8
Psychiatrically ill	7	8	4
Intervention format			
Group	19	10	10
Individual	7	3	5
Sample size, mean (SD), No.	116.1 (115.0)	130.9 (115.0)	143.3 (116.6)
Age, mean (SD), y	28.7 (13.9)	28.2 (13.7)	42.2 (16.1)
Women, mean (SD), %	58.9 (23.1)	60.2 (38.1)	72.0 (20.7)
Sessions, mean (SD), No.	8.8 (8.6)	10.0 (4.6)	5.6 (3.7)
Length of sessions, mean (SD), min	123.8 (102.3)	87.3 (24.8)	102.5 (102.8)
SASQI score, mean (SD)	10.0 (2.0)	10.9 (1.3)	10.3 (1.5)

Abbreviations: SASQI, Scale for Assessing Scientific Quality of Investigations (modified version).

<sup>a</sup> One study<sup>78</sup> included both adolescents and young adults (aged 13-25 years).

groups, 27 RCTs (47%) reported significantly improved wisdom components with a medium to large effect size.<sup>39,45,47,49-51,55-57,59,60,62-65,67,69,70,72,74,77,82-84,88,90,92</sup>

The proportion of trials with medium to large effect sizes did not differ across the 3 components: prosocial behaviors (15 of 29 studies<sup>39,45,47,49-51,55-57,59,60,62,63,65</sup>), emotional regulation (6 of 13 studies<sup>67,69,70,72,74,77</sup>), and spirituality (6 of 15 studies<sup>82-84,88,90,92</sup>). Among the prosocial behavior RCTs, the trials with medium to large effect sizes had older mean participant ages ( $t_{20} = -3.59$ ;  $P = .002$ ;  $d, -1.57$ ). Among the emotional regulation studies, the interventions with medium to large effect sizes were more likely to occur within individual-based formats (3 of 3<sup>69,70,77</sup>) compared with group settings (3 studies<sup>67,72,74</sup> of 10 studies<sup>66-68,71-76</sup>) ( $\chi^2_1 = 4.55$ ;  $P = .03$ ).

The outcome measures used varied across the reports (eTable 3 in the Supplement). Forty-eight of 57 studies<sup>41-52,54-62,65,70-75,77-92</sup> (84%) used self-rated measures. Ten trials (17.5%) used objective outcomes: 5 on prosocial behaviors (3 task-based,<sup>53,63,64</sup> 1 researcher-rated,<sup>39</sup> and 1 parent-rated<sup>40</sup>), and 5 on emotional regulation in 4 studies (teacher-rated or parent-rated).<sup>66-69</sup>

### Prosocial Behavior Interventions

Twenty-nine RCTs focused on prosocial behaviors of empathy, compassion (including self-compassion), and altruism.<sup>39-65</sup> Twenty-three studies<sup>39,43-52,55-60,62-65</sup> included adults. Eight trials<sup>40,47-49,52,55</sup> included persons with psychiatric or behavioral problems, 1 trial<sup>50</sup> included people with physical illness (diabetes mellitus), and 21 were community based.<sup>41-46,51,53,54,56-65</sup> While empathy and compassion were hypothesized to reduce cyberbullying<sup>41,54</sup> and aggression<sup>61</sup> and improve grades and learning<sup>53</sup> in young people, self-

compassion was hypothesized to decrease distress and improve well-being.<sup>50,62</sup>

The meta-analysis calculations for these interventions were based on 23 RCTs from 21 articles,<sup>39-41,43,45-47,49,50,52-62,65</sup> with a pooled SMD of 0.43 (95% CI, 0.22-0.3;  $P = .01$ ; Figure 2). Heterogeneity of the studies was considerable ( $I^2 = 84\%$ ;  $P < .001$ ). Altogether, the interventions had a statistically significant but modest association with prosocial behaviors.

### Emotional Regulation Interventions

There were 13 studies (in 12 articles<sup>66-77</sup>) of emotional regulation interventions in either psychiatrically ill or community-based populations, with 10 using group settings.<sup>66-68,71-76</sup> These RCTs sought to improve binge-eating behaviors,<sup>71</sup> behavioral problems,<sup>69</sup> test anxiety,<sup>72</sup> and family relationships.<sup>67</sup>


The meta-analysis calculations for these interventions were based on 12 studies from 11 published RCTs,<sup>66,67,69-77</sup> with a pooled SMD of 0.67 (95% CI, 0.21-1.12;  $P = .004$ ; Figure 3). Heterogeneity of study outcomes was considerable ( $I^2 = 93\%$ ;  $P < .001$ ). Altogether, the interventions had a statistically significant, medium-sized association with emotional regulation outcomes.

### Spirituality Interventions

There were 15 spirituality-focused RCTs,<sup>78-92</sup> all in adults, and 8 of these<sup>80,82-88</sup> were in people with serious and/or terminal medical illnesses. Four RCTs included psychiatric samples: patients with opiate use disorders stabilized with methadone maintenance,<sup>90</sup> adults with depression/anxiety,<sup>78,81</sup> and women with eating disorders.<sup>91</sup> Ten interventions were conducted in group settings<sup>80,81,83,84,86,87,89-92</sup> and 2 drew from specific religions (Buddhist teachings<sup>80</sup> or Islamic traditions<sup>82</sup>).



Figure 2. Forest Plot for Interventions for Prosocial Behaviors



SMD indicates standardized mean differences.

Spirituality was hypothesized to reduce psychological suffering and improve quality of life.<sup>78-92</sup>

The meta-analysis calculations for these interventions were based on 12 studies from 12 published RCTs,<sup>78-92</sup> with a pooled SMD of 1.00 (95% CI, 0.41-1.60;  $P = .001$ ; Figure 2C). Heterogeneity of study outcomes was considerable ( $I^2 = 96\%$ ;  $P < .001$ ). Altogether, the interventions had a statistically significant large-sized association with spiritual outcomes.

### Quality of Included Studies

The quality rating for each study is reported in eTable 2 in the Supplement, using the information from the modified SASQI scale. Overall, 29 studies<sup>40-43,45,46,48,50,51,54,55,57,59,60,64,65,69,71,74-76,78,81,82,85,87-90</sup> (51%) described the method used to randomize participants to the intervention vs control groups, 23 (40%) included an active control group,<sup>40,44,47,49,51,54,58,59,61,63,66,68,74-76,79,82,85,87,88,91,92</sup> and 21 (37%) examined whether participants who dropped out differed significantly from those who completed the study.<sup>40,45,46,48-51,55,56,69,73-75,77,78,82,83,85,90,92</sup> The lower-quality reports (those in the lower 50th percentile; 25 studies vs 32 higher-quality reports) were less likely to describe randomization methods (6 of 25 lower-quality studies vs 13 of 32 higher-quality studies), inclusion and exclusion criteria (14 lower-quality studies vs 30 higher-quality studies), withdrawals or dropouts (11 lower-quality studies vs 31 higher-quality studies), comparison of demographic variables between the control and intervention groups (18 lower-quality

studies vs 31 higher-quality studies), and analytical plan to address differences between control vs intervention groups (15 lower-quality studies vs 29 higher-quality studies) and between participants who dropped out vs completed the study (0 lower-quality studies vs 29 higher-quality studies).


### Publication Bias

The Egger test indicated the presence of publication bias among the studies of prosocial behavior ( $z$ , 3.48;  $P < .001$ ) and emotional regulation ( $z$ , 9.01;  $P < .001$ ), but not the spirituality studies ( $z$ , 0.10;  $P = .92$ ). Funnel plots using the trim-and-fill method<sup>36</sup> resulted in the following adjusted SMDs (prosocial behavior: SMD, 0.4 [95% CI, 0.16-0.78];  $P = .003$ ; emotional regulation: SMD, 1.26 [95% CI, -0.53 to 3.04];  $P = .17$ ; spirituality: SMD, 1.00 [95% CI, 0.46-1.54];  $P < .001$ ; eFigures 1, 2, and 3 in the Supplement).

### Meta-regression Analysis

The meta-regression analyses found that effect sizes did not differ among different wisdom components. Among trials on prosocial behavior, sample populations with older mean ages had higher effect sizes ( $\beta$ , 0.08 [SE, 0.04]); for spirituality trials, sample populations with younger mean ages had higher effect sizes ( $\beta$ , -0.13 [SE, 0.04]) (Table 2). Among spirituality interventions, studies with higher SASQI scores had larger effect sizes ( $\beta$ , 4.17 [SE, 1.07]), while the reverse was true for prosocial behavior trials ( $\beta$ , -0.91 [SE, 0.44]).

Figure 3. Forest Plots for Interventions for Emotional Regulation and Spirituality



SMD indicates standardized mean differences.

## Discussion

Overall, our meta-analysis of RCTs supports significant increase in levels of the wisdom components with small to large effect sizes, with no difference among the 3 wisdom components. However, the studies had several limitations.

The large range of SMD values could reflect differing outcome measures, as well as variability in types of study participants, control conditions, and interventions. The meta-regression analysis took into account several variables listed in Table 2. The intervention methodology varied widely from cognitive behavioral therapy to mindfulness-based group therapies to spiritual counseling. Other participant characteristics and nuances in intervention type, adherence, and outcome measures that could not be assessed in this study might also affect SMD value.

## Importance of Participant Characteristics

Prosocial behavioral interventions conducted in samples with older mean ages had greater effect sizes. This finding is consistent with published cross-sectional studies reporting higher levels of certain wisdom components in participants of older age.<sup>93-95</sup> These include emotional regulation,<sup>96-98</sup> with decreased likelihood of using destructive strategies to manage conflict<sup>99</sup> and lower recall of negative emotional experiences.<sup>100</sup> Similarly, older adults exhibit greater emotional empathy (empathic concern)<sup>101</sup> and altruistic behavior,<sup>21,102</sup> despite deficits in cognitive empathy (accurate perceptions of others' feelings).<sup>101</sup>

The percentage of female participants varied across studies, and male and female participants had similar responses to interventions. However, women have been reported to have greater baseline empathy and compassion,<sup>103</sup> as well as different emotional neurocircuitry than men.<sup>104-106</sup> One study<sup>67</sup> of emotional regulation in family units found a medium to large

Table 2. Results of Meta-regression Analysis

Characteristic	Multivariate model of all studies		
	$\beta$ (SE)	Wald statistic	P value
Intercept	-1.12 (0.80)	1.93	.17
Wisdom component			
Emotional regulation (vs prosocial behaviors)	-0.12 (0.53)	0.05	.82
Spirituality (vs prosocial behaviors)	-0.32 (0.42)	0.57	.45
Participant type			
With physical illness (vs community based)	0.59 (0.56)	1.11	.29
With psychiatric illness (vs community based)	0.09 (0.34)	0.07	.79
Mean age, y	0.08 (0.04)	5.11	.02
Women, %	0.003 (0.008)	0.16	.69
Intervention format			
Individual (vs group)	0.39 (0.42)	0.87	.35
Sessions, No.	-0.02 (0.03)	0.55	.46
Length of sessions, min	0.002 (0.002)	1.03	.31
SASQI			
Scores in top 50th percentile (vs bottom 50th percentile)	-0.91 (0.44)	4.39	.04
Interaction terms			
Wisdom component (prosocial behavior) $\times$ mean age	0.00 [Reference]	NA	NA
Wisdom component (emotional regulation) $\times$ mean age	-0.06 (0.04)	2.18	.14
Wisdom component (spirituality) $\times$ mean age	-0.13 (0.04)	9.64	.002
Wisdom component (prosocial behavior) $\times$ SASQI Score (top 50th percentile)	0.00 [Reference]	NA	NA
Wisdom component (emotional regulation) $\times$ SASQI Score (top 50th percentile)	0.78 (0.87)	0.80	.37
Wisdom component (spirituality) $\times$ SASQI Score (top 50th percentile)	4.17 (1.07)	15.1	<.001

Abbreviations: NA, not applicable; SASQI, Scale for Assessing Scientific Quality of Investigations (modified version).

effect size in mothers but not in fathers, which was attributed to greater caregiving roles taken by mothers and thus greater involvement with the intervention, which had relied heavily on home-based work. Further investigation of sex differences is needed to clarify the underlying neurobiological mechanisms of positive psychological traits.

### Features of Interventions

Intervention features, such as session length, frequency, and format, were not associated with outcomes in the meta-regression. While several interventions included home-based practice, such sessions had variable guidelines and adherence levels and therefore could not be included in the meta-regression. Informal skills practice may be essential to honing such traits and should be considered in developing future interventions for wisdom components. Of note, the comparison of trials with medium to large effect sizes with trials with small effect sizes showed that, among the emotional regulation interventions, individual-based interventions were more likely to have medium to large effect sizes than group-based interventions. This finding could partially reflect the fact that fewer RCTs used individual-based formats than group formats because of cost difference and potential gains from group-based settings (eg, social and peer support).

### Study Quality

The modified SASQI scores (lower vs higher than the 50th percentile) reflect the quality of the included studies. While study

quality did not differ among different wisdom components, less rigorous RCTs of prosocial behavior had larger effect sizes overall (ie, confounding factors could inflate differences found between the intervention and control group). Interestingly, specifically among spirituality interventions, higher-quality studies reported larger effect sizes. Interventions to improve spirituality may face the additional challenges of its broad definition, ranging from awareness of other powers or forces that influence the universe to living a spiritual life (which is more similar to religious ideals),<sup>107,108</sup> connection to personality traits,<sup>6</sup> and involvement of multiple brain regions.<sup>69</sup>

### Study Outcomes

Unsurprisingly, most of the studies used self-rated or subjective measures as outcomes. While there is great value in assessing the subjective experience of individuals for assessing personality traits such as wisdom, objective assessments should be encouraged. An example is a study by Bonvicini and colleagues<sup>39</sup> objectively assessing physician empathy based on audio recordings of patient-physician interactions.

### Future Directions

Future studies should be based on hypotheses generated from this review—for example, that (1) enhancement of wisdom components will contribute to improvement of overall wisdom, health, and well-being, and (2) older people are more likely to have improvements in wisdom components than younger persons. Exploratory analyses should include examination of



hedonic vs eudemonic well-being, interventions, sex differences in response to interventions, and validity of objective vs subjective ratings of wisdom components. Use of objective measures of wisdom components, techniques such as ecological momentary assessments, technology such as artificial intelligence, relevant biomarkers, and longer-term follow-ups are recommended.

We believe that the relevance of wisdom-associated interventions expands beyond the individual level to the societal level. Over the past 3 decades, annual rates of deaths from opioid overdose and suicides have been rising rapidly,<sup>109,110</sup> resulting in a decline in the mean American lifespan for the first time in half a century.<sup>111</sup> Loneliness and social isolation have been reported to lead to higher stress levels, as well as physical and psychiatric morbidity, including substance abuse and suicidal behavior.<sup>112-115</sup> These recent behavioral epidemics of loneliness, suicides, and opioid abuse<sup>112-115</sup> will require behavioral or psychosocial vaccines or antidotes. The strong inverse correlation between loneliness and wisdom found in a recent study<sup>15</sup> suggests that wisdom may be a remedy for behavioral toxins, such as loneliness.<sup>116</sup>

### Limitations

This review has several limitations. Improvement in individual components of wisdom is not the same as increase in overall wisdom. Studies focusing on wisdom as an entity are clearly warranted. We found only 1 RCT<sup>117</sup> with overall wisdom as an outcome measure, but it did not meet other selection criteria. However, a recently published study by Treichler and colleagues<sup>118</sup> showed increased overall wisdom using the San Diego Wisdom Scale with a 1-month group-based intervention in senior housing communities, which also reduced perceived stress and increased resilience without affecting overall well-being.

Next, search terms were limited to the specific wisdom components and thus did not include negative terms associated with lack of wisdom components (eg, impulsivity, selfishness), and may have missed some relevant studies. Outcomes were limited to the specific wisdom component, because many studies did not assess well-being or other health-associated measures. Most studies relied on self-report assessments of wisdom components, which have potential for bias because of socially desirable responses and problems with recall accuracy. Objective and technology-based measures (eg, reports by participants' close associates, ecological momentary assessments, and video game-based tasks) are warranted. Whether improvements in a wisdom component generalized to everyday life was not examined. Rationale for the study design, study participants' sociodemographic and clinical

characteristics, trial methodology, outcomes evaluated, and statistics used varied across the RCTs, contributing to significant heterogeneity among studies, publication bias, and challenges in analyzing all the study features. Follow-up periods were often short, making longer-term efficacy of the interventions unknown. Another limitation pertains to exclusion of articles that were not in the English language. Also, we could not include distal outcomes that were sometimes the goals of the original RCTs (eg, reducing binge-eating behaviors, preventing cyberbullying),<sup>41,55,81</sup> because of marked heterogeneity. Finally, only a few trials were of pragmatic type (ie, combining efficacy with effectiveness, using manualized interventions, and ensuring implementation and dissemination potential in the real world).

Few studies used neurobiological assessments. One novel study, although it did not meet our selection criteria, represents the type of research required to improve our understanding of the neurobiological implications of wisdom-associated interventions. Klimecki and colleagues<sup>119</sup> used an fMRI socio-affective video task to analyze brain activation in women after compassion training compared with memory training and reported enhanced brain activation in anterior insula and anterior midcingulate cortex (regions associated with empathy), as well as the ventral striatum, anterior cingulate, and medial orbitofrontal cortex (regions identified in putative wisdom neurocircuitry).<sup>16</sup> Adding neurobiological assessments, such as regional brain activation on fMRI, with an emotion-based task (eg, seeing happy vs angry faces) would help broaden our knowledge of the brain-based mechanisms mediating improvements in wisdom or its components. Although a wisdom pill is unlikely in the near future, the next generation of advanced and targeted neurostimulation techniques could selectively activate or inhibit neurocircuits associated with components of wisdom.

### Conclusions

Basic research is needed to better understand the neurobiology of wisdom and develop new biologically oriented, wisdom-associated interventions. Eventually, development of wisdom-enhancing interventions at societal level will become a priority, although a number of steps are required to enable development and testing of effective large-scale community-wide interventions. Balancing these 2 priorities will require political wisdom on the part of health care leaders. Increased wisdom in both individuals and communities is likely to confer broad advantages in well-being and health that would ultimately improve survival and flourishing of the society as a whole.

#### ARTICLE INFORMATION

**Accepted for Publication:** March 1, 2020.

**Published Online:** May 13, 2020.

doi:10.1001/jamapsychiatry.2020.0821

**Author Affiliations:** Department of Psychiatry, University of California San Diego, La Jolla (Lee, Bangen, Avanzino, Hou, Ramsey, Eglit, Paulus, Jeste); VA San Diego Healthcare System, San Diego,

California (Lee, Bangen, Eglit); Sam and Rose Stein Institute for Research on Aging, University of California San Diego, La Jolla (Lee, Avanzino, Hou, Eglit, Liu, Tu, Jeste); Department of Family Medicine and Public Health, University of California, San Diego, San Diego (Liu, Tu); Laureate Institute for Brain Research, Tulsa, Oklahoma (Paulus); Department of Neurosciences, University of California San Diego, La Jolla (Jeste); Department of

Psychiatry, University of California San Diego, La Jolla (Lee, Bangen, Avanzino, Hou, Ramsey, Eglit, Paulus, Jeste).

**Author Contributions:** Drs Lee and Jeste had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Lee, Bangen, Avanzino, Eglit, Paulus, Jeste.

**Acquisition, analysis, or interpretation of data:** Lee, Bangen, Avanzino, Hou, Ramsey, Eglit, Liu, Tu, Jeste.

**Drafting of the manuscript:** Lee, Avanzino, Hou, Ramsey, Tu, Paulus, Jeste.

**Critical revision of the manuscript for important intellectual content:** Lee, Bangen, Avanzino, Ramsey, Eglit, Liu, Paulus, Jeste.

**Statistical analysis:** Lee, Eglit, Liu, Tu.

**Obtained funding:** Jeste.

**Administrative, technical, or material support:** Bangen, Avanzino, Eglit, Jeste.

**Supervision:** Tu, Paulus, Jeste.

**Conflict of Interest Disclosures:** Drs Lee and Jeste reported grants from National Institutes of Health, grants from Brain and Behavior Research Foundation, and grants from Stein Institute for Research on Aging during the conduct of the study. No other disclosures were reported.

**Funding/Support:** This study was supported by the NARSAD Young Investigator grant from the Brain and Behavior Research Foundation (Dr Lee), the National Institute of Mental Health (grant K23MH119375-01 [Dr Lee] and T32 Geriatric Mental Health Program grant MH019934 [Dr Jeste]), National Institutes of Health (grant UL1TR001442 of Clinical and Translational Science Award), and the Stein Institute for Research on Aging (Dr Jeste) at the University of California San Diego.

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Disclaimer:** The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health.

## REFERENCES

- Vaillant GE. *Adaptation to Life*. Harvard University Press; 1977.
- Baltes PB, Staudinger UM. Wisdom: a metaheuristic (pragmatic) to orchestrate mind and virtue toward excellence. *Am Psychol*. 2000;55(1):122-136. doi:10.1037/0003-066X.55.1.122
- Clayton VP, Birren JE. The development of wisdom across the life-span: A reexamination of an ancient topic. In: Baltes PB, Brim OG, eds. *Life-span development and behavior*. Vol 3. Academic Press; 1980:103-135.
- Sternberg RJ. A balance theory of wisdom. *Rev Gen Psychol*. 1998;2(4):347-365. doi:10.1037/1089-2680.2.4.347
- Ardelt M. Empirical assessment of a three-dimensional wisdom scale. *Res Aging*. 2003; 25(3):275-324. doi:10.1177/0164027503025003004
- Josefsson K, Cloninger CR, Hintsanen M, Jokela M, Pulkki-Råback L, Keltikangas-Järvinen L. Associations of personality profiles with various aspects of well-being. *J Affect Disord*. 2011;133(1-2): 265-273. doi:10.1016/j.jad.2011.03.023
- Jeste DV, Ardel M, Blazer D, Kraemer HC, Vaillant G, Meeks TW. Expert consensus on characteristics of wisdom. *Gerontologist*. 2010;50(5):668-680. doi:10.1093/geront/gnq022
- Blazer DG. Religion, spirituality, and mental health. *Can J Psychiatry*. 2009;54(5):281-282. doi:10.1177/070674370905400501
- Williams PB, Mangelsdorf HH, Kontra C, Nusbaum HC, Hoekner B. The relationship between mental and somatic practices and wisdom. *PLoS One*. 2016;11(2):e0149369. doi:10.1371/journal.pone.0149369
- Grossmann I. Wisdom in context. *Perspect Psychol Sci*. 2017;12(2):233-257. doi:10.1177/1745691616672066
- Webster JD, Westerhof GJ, Bohlmeijer ET. Wisdom and mental health across the lifespan. *J Gerontol B Psychol Sci Soc Sci*. 2014;69(2):209-218. doi:10.1093/geronb/gbs121
- Ardelt M, Jeste DV. Wisdom and hard times. *J Gerontol B Psychol Sci Soc Sci*. 2018;73(8):1374-1383. doi:10.1093/geronb/gbw137
- Bergsma A, Ardel M. Self-reported wisdom and happiness: an empirical investigation. *J Happiness Stud*. 2012;13(3):481-499. doi:10.1007/s10902-011-9275-5
- Ardelt M. Antecedents and effects of wisdom in old age—a longitudinal perspective on aging well. *Res Aging*. 2000;22(4):360-394. doi:10.1177/01640275000224003
- Lee EE, Depp C, Palmer BW, et al. High prevalence and adverse health effects of loneliness in community-dwelling adults across the lifespan. *Int Psychogeriatr*. 2019;31(10):1447-1462. doi:10.1017/S1041610218002120
- Meeks TW, Jeste DV. Neurobiology of wisdom. *Arch Gen Psychiatry*. 2009;66(4):355-365. doi:10.1001/archgenpsychiatry.2009.8
- Haas LF. Phineas Gage and the science of brain localisation. *J Neurol Neurosurg Psychiatry*. 2001;71(6):761-761. doi:10.1136/jnnp.71.6.761
- Bang J, Spina S, Miller BL. Frontotemporal dementia. *Lancet*. 2015;386(10004):1672-1682. doi:10.1016/S0140-6736(15)00461-4
- Brassen S, Gamer M, Peters J, Gluth S, Büchel C. Don't look back in anger! responsiveness to missed chances in successful and unsuccessful aging. *Science*. 2012;336(6081):612-614. doi:10.1126/science.1217516
- Mather M, Canli T, English T, et al. Amygdala responses to emotionally valenced stimuli in older and younger adults. *Psychol Sci*. 2004;15(4):259-263. doi:10.1111/j.0956-7976.2004.00662.x
- Beadle JN, Sheehan AH, Dahlben B, Gutches AH. Aging, empathy, and prosociality. *J Gerontol B Psychol Sci Soc Sci*. 2015;70(2):215-224. doi:10.1093/geronb/gbt091
- Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD model. *Psychol Aging*. 2002;17(1):85-100. doi:10.1037/0882-7974.17.1.85
- Davis SW, Dennis NA, Daselaar SM, Fleck MS, Cabeza R. Que PASA? the posterior-anterior shift in aging. *Cereb Cortex*. 2008;18(5):1201-1209. doi:10.1093/cercor/bhm155
- St Jacques PL, Bessette-Symons B, Cabeza R. Functional neuroimaging studies of aging and emotion: fronto-amygdala differences during emotional perception and episodic memory. *J Int Neuropsychol Soc*. 2009;15(6):819-825. doi:10.1017/S1355617709990439
- Thomas ML, Martin AS, Eyler L, et al. Individual differences in level of wisdom are associated with brain activation during a moral decision-making task. *Brain Behav*. 2019;9(6):e01302. doi:10.1002/brb3.1302
- Amstadter AB, Moscati A, Oxon MA, Maes HH, Myers JM, Kendler KS. Personality, cognitive/psychological traits and psychiatric resilience. *Pers Individ Dif*. 2016;91:74-79. doi:10.1016/j.paid.2015.11.041
- Boardman JD, Blalock CL, Button TM. Sex differences in the heritability of resilience. *Twin Res Hum Genet*. 2008;11(1):12-27. doi:10.1375/twin.11.1.12
- Zwir I, Del-Val C, Arnedo J, et al. Three genetic-environmental networks for human personality. *Mol Psychiatry*. 2019. doi:10.1038/s41380-019-0579-x
- Adler AB, Williams J, McGurk D, Moss A, Bliese PD. Resilience training with soldiers during basic combat training. *Appl Psychol Health Well Being*. 2015;7(1):85-107. doi:10.1111/aphw.12040
- Loprinzi CE, Prasad K, Schroeder DR, Sood A. Stress Management and Resilience Training (SMART) program to decrease stress and enhance resilience among breast cancer survivors. *Clin Breast Cancer*. 2011;11(6):364-368. doi:10.1016/j.clbc.2011.06.008
- Cohen J. A power primer. *Psychol Bull*. 1992;112(1):155-159. doi:10.1037/0033-2909.112.1.155
- Jeste DV, Dunn LB, Folsom DP, Zisook D. Multimedia educational aids for improving consumer knowledge about illness management and treatment decisions. *J Psychiatr Res*. 2008;42(1):1-21. doi:10.1016/j.jpsychires.2006.10.004
- Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials. *Control Clin Trials*. 1996;17(1):1-12. doi:10.1016/0197-2456(95)00134-4
- Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0*. The Cochrane Collaboration; 2011.
- Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. doi:10.1136/bmj.315.7109.629
- Duval S, Tweedie R. Trim and fill. *Biometrics*. 2000;56(2):455-463. doi:10.1111/j.0006-341X.2000.00455.x
- Tang W, He H, Tu XM. *Applied Categorical and Count Data Analysis*. Chapman & Hall/CRC; 2012. doi:10.1201/b12123
- Seber G, Lee A. *Linear Regression Analysis*. 2nd ed. John Wiley & Sons, Inc; 2003. doi:10.1002/9780471722199
- Bonvicini KA, Perlin MJ, Bylund CL, Carroll G, Rouse RA, Goldstein MG. Impact of communication training on physician expression of empathy in patient encounters. *Patient Educ Couns*. 2009;75(1):3-10. doi:10.1016/j.pec.2008.09.007
- Solantaus T, Paavonen EJ, Toikka S, Punamäki RL. Preventive interventions in families with parental depression. *Eur Child Adolesc Psychiatry*. 2010;19(12):883-892. doi:10.1007/s00787-010-0135-3
- DeSmet A, Bastiaensens S, Van Cleemput K, et al. The efficacy of the Friendly Attac serious digital game to promote prosocial bystander behavior in cyberbullying among young adolescents. *Comput Human Behav*. 2018;78:336-347. doi:10.1016/j.chb.2017.10.011

42. Influx M, Pratt M, Masalha S, Zagoory-Sharon O, Feldman R. A social neuroscience approach to conflict resolution. *Soc Neurosci*. 2019;14(4):378-389. doi:10.1080/17470919.2018.1479983
43. Larti N, Ashouri E, Aarabi A. The effect of an empathy role-play program for operating room nursing students. *J Educ Eval Health Prof*. 2018;15:29. doi:10.3352/jeehp.2018.15.29
44. Romero-Martínez Á, Lila M, Gracia E, Moya-Albiol L. Improving empathy with motivational strategies in batterer intervention programmes. *Br J Clin Psychol*. 2019;58(2):125-139. doi:10.1111/bjc.12204
45. Sommers-Spijkerman MPJ, Trompetter HR, Schreurs KMG, Bohlmeijer ET. Compassion-focused therapy as guided self-help for enhancing public mental health. *J Consult Clin Psychol*. 2018;86(2):101-115. doi:10.1037/ccp0000268
46. Buffel du Vaure C, Lemogne C, Bunge L, et al. Promoting empathy among medical students. *J Psychosom Res*. 2017;103:102-107. doi:10.1016/j.jpsychores.2017.10.008
47. Westerhof-Evers HJ, Visser-Keizer AC, Fasotti L, et al. Effectiveness of a Treatment for Impairments in Social Cognition and Emotion Regulation (T-ScEmo) after traumatic brain injury. *J Head Trauma Rehabil*. 2017;32(5):296-307. doi:10.1097/HTR.0000000000000332
48. Ascone L, Sundag J, Schlier B, Lincoln TM. Feasibility and effects of a brief compassion-focused imagery intervention in psychotic patients with paranoid ideation. *Clin Psychol Psychother*. 2017;24(2):348-358. doi:10.1002/cpp.2003
49. de Bruin EI, van der Zwan JE, Bögels SM. A RCT comparing daily mindfulness meditations, biofeedback exercises, and daily physical exercise on attention control, executive functioning, mindful awareness, self-compassion, and worrying in stressed young adults. *Mindfulness (N Y)*. 2016;7(5):1182-1192. doi:10.1007/s12671-016-0561-5
50. Friis AM, Johnson MH, Cutfield RG, Considine NS. Kindness matters: a randomized controlled trial of a mindful self-compassion intervention improves depression, distress, and HbA1c among patients with diabetes. *Diabetes Care*. 2016;39(11):1963-1971. doi:10.2337/dc16-0416
51. Galante J, Bekkers MJ, Mitchell C, Gallacher J. Loving-kindness meditation effects on well-being and altruism. *Appl Psychol Health Well Being*. 2016;8(3):322-350. doi:10.1111/aphw.12074
52. Toole AM, Craighead LW. Brief self-compassion meditation training for body image distress in young adult women. *Body Image*. 2016;19:104-112. doi:10.1016/j.bodyim.2016.09.001
53. Flook L, Goldberg SB, Pinger L, Davidson RJ. Promoting prosocial behavior and self-regulatory skills in preschool children through a mindfulness-based Kindness Curriculum. *Dev Psychol*. 2015;51(1):44-51. doi:10.1037/a0038256
54. Garaigordobil M, Martínez-Valderrey V. Effects of Cyberprogram 2.0 on "face-to-face" bullying, cyberbullying, and empathy. *Psicothema*. 2015;27(1):45-51.
55. Kelly AC, Carter JC. Self-compassion training for binge eating disorder. *Psychol Psychother*. 2015;88(3):285-303. doi:10.1111/papt.12044
56. Lor KB, Truong JT, Ip EJ, Barnett MJ. A randomized prospective study on outcomes of an empathy intervention among second-year student pharmacists. *Am J Pharm Educ*. 2015;79(2):18. doi:10.5688/ajpe79218
57. Erogul M, Singer G, McIntyre T, Stefanov DG. Abridged mindfulness intervention to support wellness in first-year medical students. *Teach Learn Med*. 2014;26(4):350-356. doi:10.1080/10401334.2014.945025
58. Mantzios M, Wilson JC. Making concrete constructs mindful: a novel approach for developing mindfulness and self-compassion to assist weight loss. *Psychol Health*. 2014;29(4):422-441. doi:10.1080/08870446.2013.863883
59. Smeets E, Neff K, Alberts H, Peters M. Meeting suffering with kindness: effects of a brief self-compassion intervention for female college students. *J Clin Psychol*. 2014;70(9):794-807. doi:10.1002/jclp.22076
60. Yadavaia JE, Hayes SC, Vilaradaga R. Using acceptance and commitment therapy to increase self-compassion. *J Contextual Behav Sci*. 2014;3(4):248-257. doi:10.1016/j.jcbs.2014.09.002
61. Castillo R, Salguero JM, Fernández-Berrocá P, Balluerka N. Effects of an emotional intelligence intervention on aggression and empathy among adolescents. *J Adolesc*. 2013;36(5):883-892. doi:10.1016/j.adolescence.2013.07.001
62. Neff KD, Germer CK. A pilot study and randomized controlled trial of the mindful self-compassion program. *J Clin Psychol*. 2013;69(1):28-44. doi:10.1002/jclp.21923
63. Weng HY, Fox AS, Shackman AJ, et al. Compassion training alters altruism and neural responses to suffering. *Psychol Sci*. 2013;24(7):1171-1180. doi:10.1177/0956797612469537
64. Riess H, Kelley JM, Bailey RW, Dunn EJ, Phillips M. Empathy training for resident physicians. *J Gen Intern Med*. 2012;27(10):1280-1286. doi:10.1007/s11606-012-2063-z
65. Wallmark E, Safarzadeh K, Daukantaite D, Maddux RE. Promoting altruism through meditation. *Mindfulness*. 2013;4(3):223-234. doi:10.1007/s12671-012-0115-4
66. Houck CD, Barker DH, Hadley W, et al. The 1-year impact of an emotion regulation intervention on early adolescent health risk behaviors. *Health Psychol*. 2016;35(9):1036-1045. doi:10.1037/hea0000360
67. Ravindran N, Engle JM, McElwain NL, Kramer L. Fostering parents' emotion regulation through a sibling-focused experimental intervention. *J Fam Psychol*. 2015;29(3):458-468. doi:10.1037/fam0000084
68. Derella OJ, Johnston OG, Loeber R, Burke JD. CBT-enhanced emotion regulation as a mechanism of improvement for childhood irritability. *J Clin Child Adolesc Psychol*. 2019;48(sup1):S146-S154. doi:10.1080/15374416.2016.1270832
69. Weiss JA, Thomson K, Burnham Riosa P, et al. A randomized waitlist-controlled trial of cognitive behavior therapy to improve emotion regulation in children with autism. *J Child Psychol Psychiatry*. 2018;59(11):1180-1191. doi:10.1111/jcpp.12915
70. Ma Y, She Z, Siu AF-Y, Zeng X, Liu X. Effectiveness of online mindfulness-based interventions on psychological distress and the mediating role of emotion regulation. *Front Psychol*. 2018;9:2090. doi:10.3389/fpsyg.2018.02090
71. Rahmani M, Omid A, Asemi Z, Akbari H. The effect of dialectical behaviour therapy on binge eating, difficulties in emotion regulation and BMI in overweight patients with binge-eating disorder. *Ment Health Prev*. 2018;9:13-18. doi:10.1016/j.mhp.2017.11.002
72. Shahidi S, Akbari H, Zargar F. Effectiveness of mindfulness-based stress reduction on emotion regulation and test anxiety in female high school students. *J Educ Health Promot*. 2017;6:87. doi:10.4103/jehp.jehp\_98\_16
73. Spidel A, Lecomte T, Kealy D, Daigneault I. Acceptance and commitment therapy for psychosis and trauma: Improvement in psychiatric symptoms, emotion regulation, and treatment compliance following a brief group intervention. *Psychol Psychother*. 2018;91(2):248-261. doi:10.1111/papt.12159
74. Tornås S, Løvstad M, Solbakk AK, Schanke AK, Stubberud J. Goal management training combined with external cuing as a means to improve emotional regulation, psychological functioning, and quality of life in patients with acquired brain injury. *Arch Phys Med Rehabil*. 2016;97(11):1841-1852.e3. doi:10.1016/j.apmr.2016.06.014
75. Schuppert HM, Timmerman ME, Bloo J, et al. Emotion regulation training for adolescents with borderline personality disorder traits. *J Am Acad Child Adolesc Psychiatry*. 2012;51(12):1314-1323.e2. doi:10.1016/j.jaac.2012.09.002
76. Berking M, Wupperman P, Reichardt A, Pejc T, Dippel A, Znoj H. Emotion-regulation skills as a treatment target in psychotherapy. *Behav Res Ther*. 2008;46(11):1230-1237. doi:10.1016/j.brat.2008.08.005
77. Slee N, Spinhoven P, Garnefski N, Arensman E. Emotion regulation as mediator of treatment outcome in therapy for deliberate self-harm. *Clin Psychol Psychother*. 2008;15(4):205-216. doi:10.1002/cpp.577
78. Rickhi B, Kania-Richmond A, Moritz S, et al. Evaluation of a spirituality informed e-mental health tool as an intervention for major depressive disorder in adolescents and young adults—a randomized controlled pilot trial. *BMC Complement Altern Med*. 2015;15:450. doi:10.1186/s12906-015-0968-x
79. Sollgruber A, Bornemann-Cimenti H, Szilagyis IS, Sandner-Kiesling A. Spirituality in pain medicine: a randomized experiment of pain perception, heart rate and religious spiritual well-being by using a single session meditation methodology. *PLoS One*. 2018;13(9):e0203336. doi:10.1371/journal.pone.0203336
80. Chimluang J, Thanasilp S, Akkayagorn L, Upasen R, Pudtong N, Tantitrakul W. Effect of an intervention based on basic Buddhist principles on the spiritual well-being of patients with terminal cancer. *Eur J Oncol Nurs*. 2017;31:46-51. doi:10.1016/j.jejon.2017.08.005
81. Sankhe A, Dalal K, Save D, Sarve P. Evaluation of the effect of spiritual care on patients with generalized anxiety and depression. *Psychol Health Med*. 2017;22(10):1186-1191. doi:10.1080/13548506.2017.1290260
82. Sajadi M, Niazi N, Khosravi S, Yaghobi A, Rezaei M, Koenig HG. Effect of spiritual counseling on spiritual well-being in Iranian women with cancer. *Complement Ther Clin Pract*. 2018;30:79-84. doi:10.1016/j.ctcp.2017.12.011



83. Labelle LE, Lawlor-Savage L, Campbell TS, Faris P, Carlson LE. Does self-report Mindfulness-Based Stress Reduction (MBSR) on spirituality and posttraumatic growth in cancer patients? *J Posit Psychol*. 2015;10(2):153-166. doi:10.1080/17439760.2014.927902
84. Brown J, Hanson JE, Schmotzer B, Webel AR. Spirituality and optimism: a holistic approach to component-based, self-management treatment for HIV. *J Relig Health*. 2014;53(5):1317-1328. doi:10.1007/s10943-013-9722-1
85. Breitbart W, Poppito S, Rosenfeld B, et al. Pilot randomized controlled trial of individual meaning-centered psychotherapy for patients with advanced cancer. *J Clin Oncol*. 2012;30(12):1304-1309. doi:10.1200/JCO.2011.36.2517
86. Chan CHY, Chan CLW, Ng EHY, et al. Incorporating spirituality in psychosocial group intervention for women undergoing in vitro fertilization. *Psychol Psychother*. 2012;85(4):356-373. doi:10.1111/j.2044-8341.2011.02040.x
87. Chochinov HM, Kristjanson LJ, Breitbart W, et al. Effect of dignity therapy on distress and end-of-life experience in terminally ill patients. *Lancet Oncol*. 2011;12(8):753-762. doi:10.1016/S1470-2045(11)70153-X
88. Ando M, Morita T, Akechi T, Okamoto T; Japanese Task Force for Spiritual Care. Efficacy of short-term life-review interviews on the spiritual well-being of terminally ill cancer patients. *J Pain Symptom Manage*. 2010;39(6):993-1002. doi:10.1016/j.jpainsymman.2009.11.320
89. Bay PS, Ivy SS, Terry CL. The effect of spiritual retreat on nurses' spirituality. *Holist Nurs Pract*. 2010;24(3):125-133. doi:10.1097/HNP.0b013e3181dd47dd
90. Margolin A, Beitel M, Schuman-Olivier Z, Avants SK. A controlled study of a spirituality-focused intervention for increasing motivation for HIV prevention among drug users. *AIDS Educ Prev*. 2006;18(4):311-322. doi:10.1521/aeap.2006.18.4.311
91. Richards PS, Berrett ME, Hardman RK, Eggett DL. Comparative efficacy of spirituality, cognitive, and emotional support groups for treating eating disorder inpatients. *Eat Disord*. 2006;14(5):401-415. doi:10.1080/10640260600952548
92. Wachholtz AB, Pargament KI. Is spirituality a critical ingredient of meditation? comparing the effects of spiritual meditation, secular meditation, and relaxation on spiritual, psychological, cardiac, and pain outcomes. *J Behav Med*. 2005;28(4):369-384. doi:10.1007/s10865-005-9008-5
93. Grossmann I, Na J, Varum ME, Park DC, Kitayama S, Nisbett RE. Reasoning about social conflicts improves into old age. *Proc Natl Acad Sci U S A*. 2010;107(16):7246-7250. doi:10.1073/pnas.1001715107
94. Happé FG, Winner E, Brownell H. The getting of wisdom. *Dev Psychol*. 1998;34(2):358-362. doi:10.1037/0012-1649.34.2.358
95. Mickler C, Staudinger UM. Personal wisdom: validation and age-related differences of a performance measure. *Psychol Aging*. 2008;23(4):787-799. doi:10.1037/a0013928
96. Van Patten R, Lee EE, Daly R, Twamley E, Tu XM, Jeste DV. Assessment of 3-dimensional wisdom in schizophrenia: associations with neuropsychological functions and physical and mental health. *Schizophr Res*. 2019;208:360-369. doi:10.1016/j.schres.2019.01.022
97. Lee EE, Martin AS, Tu X, Palmer BW, Jeste DV. Childhood adversity and schizophrenia: the protective role of resilience in mental and physical health and metabolic markers. *J Clin Psychiatry*. 2018;79(3):17m11776. doi:10.4088/JCP.17m11776
98. Khan A, Ferrari M. Wisdom of young adults with high functioning autism in Canada and Pakistan: a cross-cultural study. *Journal of Education & Social Sciences*. 2018;6(1):3-22. doi:10.20547/jess0611806101
99. Birditt KS, Fingerman KL. Do we get better at picking our battles? age group differences in descriptions of behavioral reactions to interpersonal tensions. *J Gerontol B Psychol Sci Soc Sci*. 2005;60(3):121-128. doi:10.1093/geronb/60.3.P121
100. Charles ST, Mather M, Carstensen LL. Aging and emotional memory: the forgettable nature of negative images for older adults. *J Exp Psychol Gen*. 2003;132(2):310-324. doi:10.1037/0096-3445.132.2.310
101. Sun B, Luo Z, Zhang W, Li W, Li X. Age-related differences in affective and cognitive empathy. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn*. 2018;25(5):655-672. doi:10.1080/13825585.2017.1360835
102. Sze JA, Gyurak A, Goodkind MS, Levenson RW. Greater emotional empathy and prosocial behavior in late life. *Emotion*. 2012;12(5):1129-1140. doi:10.1037/a0025011
103. Tracy LM, Giummarra MJ. Sex differences in empathy for pain. *Psychophysiology*. 2017;54(10):1549-1558. doi:10.1111/psyp.12895
104. Wu Y, Li H, Zhou Y, et al. Sex-specific neural circuits of emotion regulation in the centromedial amygdala. *Sci Rep*. 2016;6:23112. doi:10.1038/srep23112
105. Derntl B, Finkelmeyer A, Eickhoff S, et al. Multidimensional assessment of empathic abilities. *Psychoneuroendocrinology*. 2010;35(1):67-82. doi:10.1016/j.psyneuen.2009.10.006
106. Whittle S, Yücel M, Yap MB, Allen NB. Sex differences in the neural correlates of emotion. *Biol Psychol*. 2011;87(3):319-333. doi:10.1016/j.biopsycho.2011.05.003
107. Wink P, Helson R. Practical and transcendent wisdom: their nature and some longitudinal findings. *J Adult Dev*. 1997;4(1):1-15. doi:10.1007/BF02511845
108. Perry CL, Komro KA, Jones RM, Munson K, Williams CL, Jason L. The measurement of wisdom and its relationship to adolescent substance use and problem behaviors. *J Child Adolesc Subst Abuse*. 2002;12(1):45-63. doi:10.1300/J029v12n01\_03
109. Scholl L, Seth P, Kariisa M, Wilson N, Baldwin G. Drug and opioid-involved overdose deaths—United States, 2013–2017. *MMWR Morb Mortal Wkly Rep*. 2018;67(1512):1419-1427. doi:10.15585/mmwr.mm67152e1
110. Hedegaard H, Curtin SC, Warner M. Suicide rates in the United States continue to increase. Published 2018. Accessed March 31, 2020. <https://www.cdc.gov/nchs/data/databriefs/db309.pdf>
111. Arias E, Xu JQ. United States life tables, 2017. Published 2019. Accessed March 31, 2020. [https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68\\_07-508.pdf](https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68_07-508.pdf)
112. Rudd RA, Aleshire N, Zibbell J, Gladden RM. Increases in drug and opioid overdose deaths—United States, 2000–2014. Published 2016. Accessed March 31, 2020. [https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm?s\\_cid=mm6450a3\\_w](https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm?s_cid=mm6450a3_w)
113. Devitt M. CDC data show U.S. life expectancy continues to decline. Published 2018. Accessed March 31, 2020. <https://www.aafp.org/news/health-of-the-public/2018120lifexpectdrop.html>
114. CDC. Suicide rising across the US. Published 2019. Accessed March 31, 2020. <https://www.cdc.gov/vitalsigns/suicide/index.html>
115. Cacioppo JT, Hawkley LC. Social isolation and health, with an emphasis on underlying mechanisms. *Perspect Biol Med*. 2003;46(3) (suppl):S39-S52. doi:10.1353/pbm.2003.0049
116. Jeste DV, Lee EE, Cacioppo S. Battling the modern behavioral epidemic of loneliness. *JAMA Psychiatry*. Published online March 4, 2020. doi:10.1001/jamapsychiatry.2020.0027
117. Daniels LR, Boehnlein J, McCallion P. Aging, depression, and wisdom: a pilot study of life-review intervention and PTSD treatment with two groups of Vietnam veterans. *J Gerontol Soc Work*. 2015;58(4):420-436. doi:10.1080/O1634372.2015.1013657
118. Treichler EBH, Glorioso D, Lee EE, et al. A pragmatic trial of a group intervention in senior housing communities to increase resilience. *Int Psychogeriatr*. 2020;32(2):173-182. doi:10.1017/S1041610219002096
119. Klimecki OM, Leiberg S, Lamm C, Singer T. Functional neural plasticity and associated changes in positive affect after compassion training. *Cereb Cortex*. 2013;23(7):1552-1561. doi:10.1093/cercor/bhs142